arXiv:1203.3318v1 [math.CA] 15 Mar 2012

ON NEW INEQUALITIES FOR $h-$CONVEX FUNCTIONS VIA RIEMANN-LIOUVILLE FRACTIONAL INTEGRATION

MEVLÜT TUNÇ

Abstract. In this paper, some new inequalities of the Hermite-Hadamard type for $h-$convex functions via Riemann-Liouville fractional integral are given.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function and let $a, b \in I$, with $a < b$. The following inequality;

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \tag{1.1}$$

is known in the literature as Hadamard's inequality. Both inequalities hold in the reversed direction if f is concave.

In [1], Varošanec introduced the following class of functions.

Definition 1. *Let $h : J \subset \mathbb{R} \to \mathbb{R}$ be a positive function. We say that $f : I \subset \mathbb{R} \to \mathbb{R}$ is $h-$convex function or that f belongs to the class $SX(h, I)$, if f is nonnegative and for all $x, y \in I$ and $\lambda \in (0, 1)$ we have*

$$f(\lambda x + (1-\lambda)y) \leq h(\lambda)f(x) + h(1-\lambda)f(y). \tag{1.2}$$

If the inequality in (1.2) is reversed, then f is said to be $h-$concave, i.e., $f \in SV(h, I)$.

Obviously, if $h(\lambda) = \lambda$, then all nonnegative convex functions belong to $SX(h, I)$ and all nonnegative concave functions belong to $SV(h, I)$; if $h(\lambda) = \frac{1}{\lambda}$, then $SX(h, I) = Q(I)$; if $h(\lambda) = 1$, then $SX(h, I) \supseteq P(I)$ and if $h(\lambda) = \lambda^s$, where $s \in (0, 1)$, then $SX(h, I) \supseteq K_s^2$. For some recent results for $h-$convex functions we refer to the interested reader to the papers [3]-[6].

Definition 2. *[See [2]] A function $h : J \to \mathbb{R}$ is said to be a superadditive function if*

$$h(x+y) \geq h(x) + h(y) \tag{1.3}$$

for all $x, y \in J$.

In [3], Sarıkaya et al. proved the following Hadamard type inequalities for $h-$convex functions.

2000 *Mathematics Subject Classification.* 26D15, 41A51, 26D10.

Key words and phrases. Riemann-Liouville fractional integral, $h-$convex function, Hadamard's inequality.

Theorem 1. *Let* $f \in SX(h,I)$, $a,b \in I$ *with* $a<b$ *and* $f \in L_1[a,b]$. *Then*

$$\frac{1}{2h\left(\frac{1}{2}\right)} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq [f(a)+f(b)]\int_0^1 h(\alpha)d\alpha. \tag{1.4}$$

In [16], Sarıkaya et al. proved the following Hadamard type inequalities for fractional integrals as follows.

Theorem 2. *Let* $f:[a,b]\to\mathbb{R}$ *be positive function with* $0 \leq a < b$ *and* $f \in L_1[a,b]$. *If* f *is convex function on* $[a,b]$, *then the following inequalities for fractional integrals hold:*

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}(b)+J_{b^-}^{\alpha}(a)\right] \leq \frac{f(a)+f(b)}{2} \tag{1.5}$$

with $\alpha > 0$.

Now we give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 3. *Let* $f \in L_1[a,b]$. *The Riemann-Liouville integrals* $J_{a^+}^{\alpha}f$ *and* $J_{b^-}^{\alpha}f$ *of order* $\alpha > 0$ *with* $a \geq 0$ *are defined by*

$$J_{a^+}^{\alpha}f(x) = \frac{1}{\Gamma(\alpha)}\int_a^x (x-t)^{\alpha-1}f(t)dt, \quad x>a$$

and

$$J_{b^-}^{\alpha}f(x) = \frac{1}{\Gamma(\alpha)}\int_x^b (t-x)^{\alpha-1}f(t)dt, \quad x<b$$

respectively where $\Gamma(\alpha) = \int_0^{\infty} e^{-u}u^{\alpha-1}du$. *Here is* $J_{a^+}^{0}f(x) = J_{b^-}^{0}f(x) = f(x)$.

In the case of $\alpha = 1$, the fractional integral reduces to the classical integral.

For some recent results connected with fractional integral inequalities see [8]-[15] and [16].

In [16], Sarikaya *et al.* proved a variant of the identity is established by Dragomir and Agarwal in [7, Lemma 2.1] for fractional integrals as the following.

Lemma 1. *Let* $f:[a,b]\to\mathbb{R}$ *be a differentiable mapping on* (a,b) *with* $a<b$. *If* $f' \in L[a,b]$, *then the following equality for fractional integrals holds:*

$$\begin{aligned} &\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}f(b)+J_{b^-}^{\alpha}f(a)\right] \\ = &\frac{b-a}{2}\int_0^1 \left[(1-t)^{\alpha}-t^{\alpha}\right]f'(ta+(1-t)b)dt. \end{aligned}$$

The aim of this paper is to establish Hadamard type inequalities for $h-$convex functions via Riemann-Liouville fractional integral.

2. MAIN RESULTS

Theorem 3. *Let* $f \in SX(h,I)$, $a,b \in I$ *with* $a<b$ *and* $f \in L_1[a,b]$. *Then one has inequality for* $h-$*convex functions via fractional integrals*

$$\begin{aligned} &\frac{\Gamma(\alpha)}{(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}(b)+J_{b^-}^{\alpha}(a)\right] \\ \leq\ & [f(a)+f(b)]\int_0^1 t^{\alpha-1}\left[h(t)+h(1-t)\right]dt \\ \leq\ & \frac{2\left[f(a)+f(b)\right]}{(\alpha p-p+1)^{\frac{1}{p}}}\left(\int_0^1 (h(t))^q\,dt\right)^{\frac{1}{q}} \end{aligned} \tag{2.1}$$

where $p^{-1}+q^{-1}=1$.

Proof. Since $f \in SX(h,I)$, we have

$$f(tx+(1-t)y) \leq h(t)f(x)+h(1-t)f(y)$$

and

$$f((1-t)x+ty) \leq h(1-t)f(x)+h(t)f(y).$$

By adding these inequalities we get

$$f(tx+(1-t)y)+f((1-t)x+ty) \leq \left[h(t)+h(1-t)\right]\left[f(x)+f(y)\right]. \tag{2.2}$$

By using (2.2) with $x=a$ and $y=b$ we have

$$f(ta+(1-t)b)+f((1-t)a+tb) \leq \left[h(t)+h(1-t)\right]\left[f(a)+f(b)\right]. \tag{2.3}$$

Then multiplying both sides of (2.3) by $t^{\alpha-1}$ and integrating the resulting inequality with respect to t over $[0,1]$, we get

$$\begin{aligned} &\int_0^1 t^{\alpha-1}\left[f(ta+(1-t)b)+f((1-t)a+tb)\right]dt \\ \leq\ & \int_0^1 t^{\alpha-1}\left[h(t)+h(1-t)\right]\left[f(a)+f(b)\right]dt, \end{aligned}$$

$$\begin{aligned} &\frac{\Gamma(\alpha)}{(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}(b)+J_{b^-}^{\alpha}(a)\right] \\ \leq\ & [f(a)+f(b)]\int_0^1 t^{\alpha-1}\left[h(t)+h(1-t)\right]dt \end{aligned} \tag{2.4}$$

and thus the first inequality is proved.

To obtain the second inequality in (2.1), by using Hölder inequality for the right hand side of (2.4), we obtain

$$\begin{aligned}
& \int_0^1 t^{\alpha-1}\left[h(t)+h(1-t)\right]dt \\
\leq & \left(\int_0^1 \left(t^{\alpha-1}\right)^p dt\right)^{\frac{1}{p}} \left(\int_0^1 \left(h(t)+h(1-t)\right)^q dt\right)^{\frac{1}{q}} \\
= & \left(\left.\frac{t^{\alpha p-p+1}}{\alpha p-p+1}\right|_0^1\right)^{\frac{1}{p}} \left(\int_0^1 \left(h(t)+h(1-t)\right)^q dt\right)^{\frac{1}{q}} \\
= & \left(\frac{1}{\alpha p-p+1}\right)^{\frac{1}{p}} \left(\int_0^1 \left(h(t)+h(1-t)\right)^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

Then using Minkowski inequality

$$\begin{aligned}
& \left(\frac{1}{\alpha p-p+1}\right)^{\frac{1}{p}} \left(\int_0^1 \left(h(t)+h(1-t)\right)^q dt\right)^{\frac{1}{q}} \\
\leq & \left(\frac{1}{\alpha p-p+1}\right)^{\frac{1}{p}} \left[\left(\int_0^1 \left(h(t)\right)^q dt\right)^{\frac{1}{q}} + \left(\int_0^1 \left(h(1-t)\right)^q dt\right)^{\frac{1}{q}}\right] \\
= & \frac{2}{\left(\alpha p-p+1\right)^{\frac{1}{p}}} \left(\int_0^1 \left(h(t)\right)^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

where the proof is completed. □

Remark 1. *If we choose $\alpha = 1$ in Theorem 1, we obtain*

$$\begin{aligned}
\frac{1}{b-a}\int_a^b f(x)\,dx & \leq \left[f(a)+f(b)\right]\int_0^1 h(t)dt \\
& \leq \left[f(a)+f(b)\right]\left(\int_0^1 \left(h(t)\right)^q dt\right)^{\frac{1}{q}}.
\end{aligned}$$

Corollary 1. *(1) If we choose $h(\lambda) = \lambda$ in Remark 1, we get*

$$\frac{1}{b-a}\int_a^b f(x)\,dx \leq \frac{f(a)+f(b)}{2} = \frac{f(a)+f(b)}{(q+1)^{\frac{1}{q}}}$$

for ordinary convex functions.

(2) If we choose $h(\lambda) = 1$ in Remark 1, we get

$$\frac{2}{b-a}\int_a^b f(x)\,dx \leq 2\left(f(a)+f(b)\right)$$

for $P-$functions. This inequality is refinement of right hand side of (1.1) for $P-$functions.

(3) If we choose $h(\lambda) = \lambda^s$ in Remark 1, we get

$$\frac{1}{b-a}\int_0^1 f(x)\,dx \leq \frac{f(a)+f(b)}{s+1} \leq \frac{f(a)+f(b)}{(sq+1)^{\frac{1}{q}}}$$

for $s-$convex functions in the second sense with $s \in (0,1]$.

Theorem 4. *Let $f\in SX(h,I)$, $a,b\in I$ with $a<b$, h is superadditive on I and $f\in L_1[a,b]$, $h\in L_1[0,1]$. Then one has inequality for $h-$convex functions via fractional integrals*

$$\frac{\Gamma(\alpha)}{(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}(b)+J_{b^-}^{\alpha}(a)\right]\leq\frac{h(1)}{\alpha}\left[f(a)+f(b)\right]. \tag{2.5}$$

Proof. Since $f\in SX(h,I)$, we have

$$f(tx+(1-t)y)\leq h(t)f(x)+h(1-t)f(y)$$

and

$$f((1-t)x+ty)\leq h(1-t)f(x)+h(t)f(y).$$

By adding these inequalities we get

$$f(tx+(1-t)y)+f((1-t)x+ty)\leq\left[h(t)+h(1-t)\right]\left[f(x)+f(y)\right]. \tag{2.6}$$

By using (2.6) with $x=a$ and $y=b$ and h is superadditive, we get

$$f(ta+(1-t)b)+f((1-t)a+tb)\leq h(1)\left[f(a)+f(b)\right]. \tag{2.7}$$

Then multiplying both sides of (2.7) by $t^{\alpha-1}$ and integrating the resulting inequality with respect to t over $[0,1]$, we get

$$\begin{aligned}&\int_0^1 t^{\alpha-1}\left[f(ta+(1-t)b)+f((1-t)a+tb)\right]dt\\ \leq\ &\int_0^1 t^{\alpha-1}h(1)\left[f(a)+f(b)\right]dt,\end{aligned}$$

$$\begin{aligned}&\frac{\Gamma(\alpha)}{(b-a)^{\alpha}}\left[J_{a^+}^{\alpha}(b)+J_{b^-}^{\alpha}(a)\right]\\ \leq\ & h(1)\left[f(a)+f(b)\right]\int_0^1 t^{\alpha-1}dt.\end{aligned}$$

This completes the proof. □

Remark 2. *If we choose $\alpha=1$ in Theorem 4, then (2.5) reduce to special version of right hand side of (1.4).*

Theorem 5. *Let $h:J\subset\mathbb{R}\to\mathbb{R}$ and $f:[a,b]\to\mathbb{R}$ be positive functions with $0\leq a<b$ and $h^q\in L_1[0,1]$, $f\in L_1[a,b]$. If $|f'|$ is an $h-$convex mapping on $[a,b]$, then the following inequality for fractional integrals holds,*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a+}^{\alpha}f(b)+J_{b-}^{\alpha}f(a)\right]\right| \\ \leq\ &\frac{(b-a)\left[|f'(a)|+|f'(b)|\right]}{2}\left[\left(\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right)^{\frac{1}{p}}-\left(\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right)^{\frac{1}{p}}\right]\\ &\times\left[\left(\int_0^{\frac{1}{2}}(h(t))^q\,dt\right)^{\frac{1}{q}}+\left(\int_{\frac{1}{2}}^1(h(t))^q\,dt\right)^{\frac{1}{q}}\right]\end{aligned} \tag{2.8}$$

where $\alpha>0$, $p>1$ and $p^{-1}+q^{-1}=1$.

Proof. From Lemma 1 and using the properties of modulus, we have

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a+}^{\alpha}f(b)+J_{b-}^{\alpha}f(a)\right]\right| \\
\leq\ & \frac{b-a}{2}\int_0^1\left|(1-t)^{\alpha}-t^{\alpha}\right|\left|f'(ta+(1-t)b)\right|dt.
\end{aligned}$$

Since $|f'|$ is $h-$convex on $[a,b]$, we have

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a+}^{\alpha}(b)+J_{b-}^{\alpha}(a)\right]\right| \\
\leq\ & \frac{b-a}{2}\left\{\int_0^{\frac{1}{2}}\left[(1-t)^{\alpha}-t^{\alpha}\right]\left[h(t)\left|f'(a)\right|+h(1-t)\left|f'(b)\right|\right]dt\right. \\
&\left.+\int_{\frac{1}{2}}^1\left[t^{\alpha}-(1-t)^{\alpha}\right]\left[h(t)\left|f'(a)\right|+h(1-t)\left|f'(b)\right|\right]dt\right\} \\
=\ & \frac{b-a}{2}\left\{\left|f'(a)\right|\int_0^{\frac{1}{2}}(1-t)^{\alpha}h(t)\,dt-\left|f'(a)\right|\int_0^{\frac{1}{2}}t^{\alpha}h(t)\,dt\right. \\
&+\left|f'(b)\right|\int_0^{\frac{1}{2}}(1-t)^{\alpha}h(1-t)\,dt-\left|f'(b)\right|\int_0^{\frac{1}{2}}t^{\alpha}h(1-t)\,dt \\
&+\left|f'(a)\right|\int_{\frac{1}{2}}^1 t^{\alpha}h(t)\,dt-\left|f'(a)\right|\int_{\frac{1}{2}}^1(1-t)^{\alpha}h(t)\,dt \\
&\left.+\left|f'(b)\right|\int_{\frac{1}{2}}^1 t^{\alpha}h(1-t)\,dt-\left|f'(b)\right|\int_{\frac{1}{2}}^1(1-t)^{\alpha}h(1-t)\,dt\right\}.
\end{aligned} \tag{2.9}$$

In the right hand side of above inequality by using Hölder inequality for $p^{-1}+q^{-1}=1$ and $p>1$, we get

$$\int_0^{\frac{1}{2}}(1-t)^{\alpha}h(t)\,dt=\int_{\frac{1}{2}}^1 t^{\alpha}h(1-t)\,dt\leq\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}},$$

$$\int_0^{\frac{1}{2}}(1-t)^{\alpha}h(1-t)\,dt=\int_{\frac{1}{2}}^1 t^{\alpha}h(t)\,dt\leq\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\left(\int_{\frac{1}{2}}^1[h(t)]^q\,dt\right)^{\frac{1}{q}},$$

$$\int_0^{\frac{1}{2}}t^{\alpha}h(t)\,dt=\int_{\frac{1}{2}}^1(1-t)^{\alpha}h(1-t)\,dt\leq\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}}$$

and

$$\int_0^{\frac{1}{2}}t^{\alpha}h(1-t)\,dt=\int_{\frac{1}{2}}^1(1-t)^{\alpha}h(t)\,dt\leq\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\left(\int_{\frac{1}{2}}^1[h(t)]^q\,dt\right)^{\frac{1}{q}}.$$

Then using the above inequalities in the right hand side of (2.9), we get

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}}\left[J_{a+}^{\alpha}(b)+J_{b-}^{\alpha}(a)\right]\right| \\
\leq\ & \frac{b-a}{2}\left\{|f'(a)|\left\{\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}}\right\}\right. \\
&+\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left(\int_{\frac{1}{2}}^{1}[h(t)]^q\,dt\right)^{\frac{1}{q}}\Bigg\} \\
&+|f'(b)|\left\{\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left(\int_{\frac{1}{2}}^{1}[h(t)]^q\,dt\right)^{\frac{1}{q}}\right. \\
&+\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}}\Bigg\} \\
=\ & \frac{b-a}{2}\left\{|f'(a)|\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left[\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}}+\left(\int_{\frac{1}{2}}^{1}[h(t)]^q\,dt\right)^{\frac{1}{q}}\right]\right. \\
&+|f'(b)|\left(\left[\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}-\left[\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right]^{\frac{1}{p}}\right)\left[\left(\int_0^{\frac{1}{2}}[h(t)]^q\,dt\right)^{\frac{1}{q}}+\left(\int_{\frac{1}{2}}^{1}[h(t)]^q\,dt\right)^{\frac{1}{q}}\right]\Bigg\} \\
=\ & \frac{(b-a)\left[|f'(a)|+|f'(b)|\right]}{2}\left[\left(\frac{2^{\alpha p+1}-1}{2^{\alpha p+1}(\alpha p+1)}\right)^{\frac{1}{p}}-\left(\frac{1}{2^{\alpha p+1}(\alpha p+1)}\right)^{\frac{1}{p}}\right] \\
&\times\left[\left(\int_0^{\frac{1}{2}}(h(t))^q\,dt\right)^{\frac{1}{q}}+\left(\int_{\frac{1}{2}}^{1}(h(t))^q\,dt\right)^{\frac{1}{q}}\right]
\end{aligned}$$

which is the desired result. The proof is completed. □

REFERENCES

[1] S. Varošanec, *On $h-$convexity*, J. Math. Anal. Appl., 326 (2007), 303-311.

[2] H. Alzer, A superadditive property of Hadamard's gamma function, Abh. Math. Semin. Univ. Hambg., 79 (2009), 11-23.

[3] M. Z. Sarıkaya, A. Sağlam and H. Yıldırım, On some Hadamard-type inequalities for h-convex functions. Journal of Mathematical Inequalities, 2 (3) (2008), 335-341.

[4] M. Bombardelli and S. Varošanec, Properties of $h-$convex functions related to the Hermite-Hadamard-Fejér inequalities, Computers and Mathematics with Applications, 58 (2009), 1869-1877.

[5] M.Z. Sarıkaya, E. Set and M.E. Özdemir, On some new inequalities of Hadamard type involving $h-$convex functions, Acta Math. Univ. Comenianae, Vol. LXXIX, 2 (2010), pp. 265-272.

[6] P. Burai and A. Házy, On approximately $h-$convex functions, Journal of Convex Analysis, 18 (2) (2011).

[7] S.S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and trapezoidal formula, Appl. Math. Lett., 11 (5) (1998), 91-95.

[8] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.

[9] Z. Dahmani, *New inequalities in fractional integrals*, International Journal of Nonlinear Science, 9(4) (2010), 493-497.

[10] M.E. Özdemir, H. Kavurmacı and M. Avcı, New inequalities of Ostrowski type for mappings whose derivatives are (α, m)-convex via fractional integrals, RGMIA Research Report Collection, 15(2012), Article 10, 8 pp.

[11] M.E. Özdemir, H. Kavurmacı and Ç. Yıldız, Fractional integral inequalities via s-convex functions, arXiv:1201.4915v1 [math.CA] 24 Jan 2012.

[12] E. Set, New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals, Comput. Math. Appl., In Press, Corrected Proof, 29 December 2011.

[13] Z. Dahmani, *On Minkowski and Hermite-Hadamard integral inequalities via fractional integration*, Ann. Funct. Anal. 1(1) (2010), 51-58.

[14] Z. Dahmani and L. Tabharit, S. Taf, *Some fractional integral inequalities*, Nonl. Sci. Lett. A., 1(2) (2010), 155-160.

[15] Z. Dahmani, L. Tabharit and S. Taf, *New generalizations of Grüss inequality using Riemann-Liouville fractional integrals*, Bull. Math. Anal. Appl., 2(3) (2010), 93-99.

[16] M. Z. Sarıkaya, E. Set, H. Yaldiz and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Mathematical and Computer Modelling, In Press.

University of Kilis 7 Aralik, Faculty of Science and Arts, Department of Mathematics, 79000, Kilis, Turkey
E-mail address: `mevluttunc@kilis.edu.tr`